

DOI: 10.14744/JPD.2024.9_237 J Pediatr Dent 2025;11(1):9-16



Original Article

Nigerian Parents' Awareness and Practices on Tetracycline Use in Children and Oral Side Effects

Nneka Maureen Chukwumah¹ Joan Emien Enabulele²

¹Department of Pedodontics, School of Dentistry, College of Medical Sciences, University of Benin, Benin City, Nigeria

²Department of Restorative Dentistry, School of Dentistry, College of Medical Sciences, University of Benin, Benin City, Nigeria

Address for correspondence: Nneka Maureen Chukwumah, Department of Pedodontics, School of Dentistry, College of Medical Sciences, University of Benin, Edo State, Nigeria **E-mail:** nneka.chukwumah@uniben.edu

Abstract

Objective: Misuse of antibiotics, including tetracycline, is prevalent in developing countries, posing risks to children's oral health. This study aimed to explore tetracycline use in children and parental awareness and practice of its oral side effects.

Materials and Methods: A cross-sectional study involving 267 parents/caregivers was conducted. Participants completed a semi-structured questionnaire that collected data on socio-demographic characteristics, knowledge, attitude, and practice of antibiotic use, as well as knowledge of the oral implications of tetracycline use. Data were analyzed to identify associations between socio-demographic characteristics, and awareness of oral adverse effects.

Results: A total of 267 respondents participated in the study. Among them, 23.6% were aged 31–35 years, 71.5% were female, 78.3% had a tertiary level of education, 88.0% were married, and 48.3% were skilled workers. About 39.3% of respondents reported using tetracycline in children, with most usage being self-prescribed (64.2%). Only 122 respondents (45.7%) received information about tetracycline use, with friends and family being the primary information source (37.2%). While the majority of respondents (62.2%) knew that using tetracycline in children led to adverse events, less than half cited tooth discoloration (34.1%) as an adverse effect. Significant associations were found between the level of education (p=0.007), occupation (p=0.009), and tetracycline use.

Conclusion: This study highlights poor knowledge, practice and misuse of antibiotics and tetracycline in children, with inadequate parental/caregiver awareness of oral side effects. Education and awareness campaigns are crucial to promote responsible antibiotic use, focusing on appropriate antibiotic use, potential oral adverse effects and medical supervision.

Keywords: Adverse effects, children, knowledge, parents, tetracycline-use

Introduction

Tetracycline is a broad-spectrum antibiotic used to treat various infections, but its use in children is cautious due to risks of permanent tooth stains and other side effects. [1,2] Pregnancy and childhood are critical periods where tetracycline use is approached with caution due to its potential to cause permanent tooth discoloration and other adverse effects.[1,2] Tetracycline use during tooth development can cause irreversible discoloration.[3–7] Misuse and overuse of antibiotics are prevalent in developing countries, accelerating antimicrobial resistance and limiting treatment options.[8,9] In Nigeria, self-medication and misuse of antibiotics are common, and parents lack knowledge about proper use and potential adverse effects in children, contributing to public health risks.[10]

How to cite this article: Chukwumah NM, Enabulele JE. Nigerian Parents' Awareness and Practices on Tetracycline Use in Children and Oral Side Effects. J Pediatr Dent 2025;11(1):9-16



The American Academy of Pediatrics recommends reserving tetracycline use in children for infections where benefits outweigh risks.[11] Tetracycline use during tooth development can cause irreversible discoloration, leading to emotional distress and psychosocial problems.[3,12] Educating patients and the public about proper medication adherence is crucial to mitigate these risks.[12] This study aims to explore tetracycline use in children and parental awareness of its oral effects, addressing a critical knowledge gap.[5-7,12] The findings will inform parent education strategies and policymakers, promoting responsible use and mitigating adverse health consequences, emphasizing the risks of antibiotic misuse and potential harm to children's health.[13] Additionally, this study aims to inform policymakers, providing insights for stricter regulations on prescription-only medications, promoting responsible use, and mitigating adverse health consequences.

By investigating tetracycline use and awareness, this study seeks to contribute meaningfully to improving public health outcomes in Nigeria and beyond. The alarming rate of antibiotic misuse in Nigeria justifies this study, highlighting the need for targeted education and stricter regulations to promote responsible antibiotic use and mitigate adverse health consequences.[14,15]

Materials and Methods

Study design

This was a cross-sectional study. It investigated the use of tetracycline in children and parental/caregiver awareness of its oral side effects.

Study participants

Adults accompanying children to the dental outpatient clinic at the University of Benin Teaching Hospital were recruited.

Sample size

Using Cochran's formula, a sample size of 241 was calculated, based on a 95% confidence level, 5% error margin, and 19.5% prevalence of awareness from a previous study.[16,17] The calculated sample size was adjusted to account for attrition, giving a total sample size of 267.

Sampling procedure

Parents who brought their children to the pediatric dental clinic and provided informed consent were included.

Data collection

A self-administered, semi-structured questionnaire gathered data on:

- 1. Socio-demographic characteristics
- 2. Knowledge of antibiotics and usage
- 3. Attitude and practice of antibiotic use
- 4. Knowledge of oral implications of tetracycline use

Data analysis

IBM SPSS version 26.0 was used for analysis. Descriptive statistics summarized socio-demographic characteristics. Inferential statistics, such as the chi-square test, were employed to examine associations between socio-demographic characteristics and tetracycline use, knowledge of adverse effects, and awareness of oral adverse effects. Results were presented in tables and charts, with a significance level of p<0.05 and a 95% confidence interval.

Ethical considerations

Ethical approval was obtained from the University of Benin Teaching Hospital Ethical and Research Committee. Informed consent was obtained from participants, ensuring confidentiality and anonymity.

Results

Socio-demographics

A total of 267 Nigerian parents responded to the survey, with a mean age of 40.48 ± 8.6 years. The most represented age group was 31-35 years (23.6%). The majority of participants were female (71.5%), married (88.0%), and had a tertiary level of education (78.3%), while skilled workers comprised 48.3% (Table 1).

Tetracycline use and perceptions

Only 42.7% believed tetracycline could cure infections in children, while 51.7% thought it could treat dysentery and diarrhea; however, 19.5% disagreed. A total of 60.7% thought children could abuse tetracycline, and 39.3% reported using tetracycline in children, with 64.2% not obtaining a doctor's prescription. The frequency and duration of use varied widely (Figs. 1, 2). There was a significant association between the level of education and tetracycline use (p=0.007) and between occupation and tetracycline use (p=0.009). However, there were no significant associations with age, gender, or marital status and tetracycline use. Specifically, a higher proportion of unskilled workers (55.6%) and semi-skilled workers (50.0%) reported using tetracycline in children (Table 2).

Reasons for tetracycline use

The primary reason for tetracycline use in children was to treat dysentery and diarrhea (83.6%), while the least

Characteristics	Frequency	Percent	CI
Age group			
≤30 years	22	8.2	5.2–11.6
31–35 years	63	23.6	18.7–28.5
36–40 years	59	22.1	17.2–27.3
41–45 years	59	22.1	17.2–27.3
46–50 years	39	14.6	10.5–19.1
>50 years	25	9.4	6.0–12.7
Gender			
Male	76	28.5	23.2-34.5
Female	191	71.5	65.5–76.8
Level of education	-		
Primary	14	5.2	2.6-8.2
Secondary	44	16.5	12.0–21.0
Tertiary	209	78.3	73.0-83.1
Marital status			
Single	32	12.0	8.2–16.1
Married	235	88.0	83.9–91.8
Occupation			
Professional	47	17.6	13.1–22.1
Skilled worker	129	48.3	42.7–54.3
Semi-skilled worker	18	6.7	4.1–10.1
Unskilled worker	63	23.6	18.7–28.5
Dependent	10	3.7	1.5–6.4
Total	267	100.0	

Table 1. Sociodemographic characteristics of the respondents

CI: Confidence interval

common reason was to treat cough (3.7%). There was a significant association between the level of education and reasons for use (p=0.05), showing a decrease in tetracycline use as the level of education increased.

Information sources

There was poor knowledge among respondents regarding the dose and duration of tetracycline (Figs. 1, 2). The primary sources of information and prescriptions for tetracycline use were friends and family (37.2%), doctors (26.2%), and pharmacists (24.1%).

Adverse effects and awareness of oral adverse effects

Only 62.2% believed tetracycline use could cause adverse events, and the most commonly cited were tooth discoloration (34.1%), nausea (25.5%), and vomiting (16.5%). Additionally, only 37.8% were aware of oral adverse effects. A statistically significant association was documented between the level of education and awareness (p=0.01) and between occupation and awareness (p=0.001) of tetracycline use (Table 3).





Safety perceptions

Concerning the safety of tetracyclines, 41.9% were unsure about their safety, 24.3% were confident in their safety, and 33.7% were certain they were not safe for children. The majority (63.7%) believed tetracycline was not safe for use during pregnancy, while 97.4% had never advised a pregnant woman to use it.

Characteristics	Use	of tetracyc	line in chilo	dren	Total	
	Ye	Yes		lo		
	n	%	n	%	n	%
Level of education					p=0	.007
Primary	9	64.3	5	35.7	14	100.0
Secondary	24	54.5	20	45.5	44	100.0
Tertiary	72	34.4	137	65.6	209	100.0
Age group					p=0).53
≤30 years	9	40.9	13	59.1	22	100.0
31–35 years	19	30.2	44	69.8	63	100.0
36–40 years	25	42.4	34	57.6	59	100.0
41–45 years	22	37.3	37	62.7	59	100.0
46–50 years	18	46.2	21	53.8	39	100.0
>50 years	12	48.0	13	52.0	25	100.0
Gender					p=0).25
Male	34	44.7	42	55.3	76	100.0
Female	71	37.2	120	62.8	191	100.0
Occupation					p=0	.009
Professional	11	23.4	36	76.6	47	100.0
Skilled worker	47	36.4	82	63.6	129	100.0
Semi-skilled worker	9	50.0	9	50.0	18	100.0
Unskilled worker	35	55.6	28	44.4	63	100.0
Dependent	3	30.0	7	70.0	10	100.0
Marital status					p=0	0.35
Single	15	46.9	17	53.1	32	100.0
Married	90	38.3	145	61.7	235	100.0
Total	105	39.3	162	60.7	267	100.0

Table 2. Association between socio-demographic characteristics and use of tetracycline in children

Discussion

This study investigated Nigerian parents' awareness and practices regarding tetracycline use in children and its oral side effects. The findings revealed a limited understanding of tetracycline's effectiveness in treating childhood infections, with many parents overestimating its capabilities. Concerns about misuse and adverse effects were prevalent, with tooth discoloration, nausea, and vomiting being the most commonly mentioned side effects. The study highlighted the need for education and awareness campaigns to promote responsible antibiotic use, improve knowledge about potential oral adverse effects, and reduce the risk of antibiotic resistance.

The sociodemographic characteristics of the respondents showed parents between 24-76 years, with a mean age of 40.48±8.6 years. The 31–35-year-old age group formed the bulk of parents seeking dental care for their children, and females comprised 71.5% of the respondents, 88% of whom were married. This is consistent with previous reports,[18–20] which also state that mothers are more likely to present with their children.[21] This finding aligns with the views of researchers who argue that mothers and women can play a vital role in maintaining children's oral health if they possess the necessary knowledge.[22] Thus, it may be important to target females and mothers during oral health awareness or promotion campaigns.

At least 78.3% of this study population had a tertiary level of education. Level of education and occupation are significant predictors of awareness about oral adverse effects. This study found a significant association

Table 3. Association betw	veen sociode	mographic cł	naracteristics	, knowledge	of side effec	ts and tetrac	ycline use in	children				
Characteristics	Presence	of side effect chilc	s of tetracyc Iren	line use in	Ď	tal	Knowled	ge of tetrac	ycline use in	children	To	tal
	*	les	Z	0			3,	SS	Z	0		
	٢	%	ء	%	٢	%	۲	%	E	%	5	%
Level of education					b=0	.29)=d	.01
Primary	∞	57.1	9	42.9	14	100.0	£	21.4	11	78.6	14	100.0
Secondary	23	52.3	21	47.7	44	100.0	6	20.5	35	79.5	44	100.0
Tertiary	135	64.6	74	35.4	209	100.0	89	42.6	120	57.4	209	100.0
Age group					b=0	.01					b=0	007
≤30 years	15	68.2	7	31.8	22	100.0	10	45.5	12	54.5	22	100.0
31–35 years	47	74.6	16	25.4	63	100.0	31	49.2	32	50.8	63	100.0
36–40 years	36	61.0	23	39.0	59	100.0	23	39.0	36	61.0	59	100.0
41–45 years	25	42.4	34	57.6	59	100.0	10	16.9	49	83.1	59	100.0
46–50 years	26	66.7	13	33.3	39	100.0	15	38.5	24	61.5	39	100.0
>50 years	17	68.0	8	32.0	25	100.0	12	48.0	13	52.0	25	100.0
Gender					b=C	.83)=d	.73
Male	48	63.2	28	36.8	76	100.0	30	39.5	46	60.5	76	100.0
Female	118	61.8	73	38.2	191	100.0	71	37.2	120	62.8	191	100.0
Occupation					b=C	.10					b=0	.001
Professional	35	74.5	12	25.5	47	100.0	29	61.7	18	38.3	47	100.0
Skilled worker	80	62.0	49	38.0	129	100.0	46	35.7	83	64.3	129	100.0
Semi-skilled worker	7	38.9	11	61.1	18	100.0	2	11.1	16	88.9	18	100.0
Unskilled worker	39	61.9	24	38.1	63	100.0	19	30.2	44	69.8	63	100.0
Dependent	5	50.0	5	50.0	10	100.0	5	50.0	5	50.0	10	100.0
Marital status					b=0	.02)=d	.54
Single	26	81.3	6	18.8	32	100.0	17	53.1	15	46.9	32	100.0
Married	140	59.6	95	40.4	235	100.0	84	35.7	151	64.3	235	100.0
Total	166	62.2	101	37.8	267	100.0	101	35.7	162	60.7	267	100.0

between the level of education and awareness of oral adverse effects, with individuals with higher education levels possessing greater knowledge, which is comparable with a previous report[20] and also supported by other research stating that education is a significant predictor of health-seeking behavior.[23,24] The occupation distribution, comprising mostly skilled workers (48.3%), professionals (17.6%), and unskilled workers (23.6%), provides insight into the respondents' socioeconomic status. This is in line with studies highlighting occupation as a determinant of health outcomes. [25–27] Occupation also played a crucial role, with healthcare professionals being more likely to be aware of potential oral adverse effects due to their occupationspecific education and training.[17,28]

With these results, comprising mostly females, married individuals, and participants with tertiary levels of education, one would expect a higher level of awareness regarding antibiotic use. However, the results indicate significant misconceptions about tetracycline effectiveness and safety. These findings have important implications for public health education and awareness campaigns, which should prioritize targeting individuals at all levels of education, both professionals and non-professionals, to improve knowledge about antibiotic use and its potential consequences.

Respondents demonstrated a limited understanding of tetracycline's effectiveness in treating childhood infections, with 83.6% indicating its use for treating dysentery and diarrhea. This is consistent with previous findings,[17] indicating a significant knowledge gap. This ambiguity may stem from inadequate knowledge or misinformation about appropriate antibiotic use.[29,30] This finding suggests that parents may rely on tetracyclines as a broad-spectrum solution rather than understanding its specific application. These misconceptions and lack of knowledge about antibiotics can lead to inappropriate use,[31] with 60.7% of respondents agreeing that self-prescription and misuse are possible, which could contribute to antibiotic resistance.

The findings emphasize the need for interventions and education to improve knowledge and understanding of antibiotic use among parents and caregivers, especially regarding children, ensuring appropriate use and reducing the risk of antibiotic resistance. [19] Another disturbing trend was friends and family (37.2%) being the primary source of information for tetracycline use, followed by doctors and pharmacists (26.2% and 24.1%, respectively). A significant proportion of respondents (39.3%) had used tetracycline in children, which is particularly concerning given the potential risks associated with antibiotic misuse and its restriction for use in children under 8 years of age due to the risk of permanent tooth discoloration.[32] Furthermore, the majority (64.2%) of those who used tetracycline in children did so without a doctor's prescription, highlighting a dangerous trend and a potential lack of medical supervision and guidance.[11] The variability in daily dosage and duration of tetracycline use among respondents, as reported in a previous study,[17] is also noteworthy. Healthcare professionals should emphasize the importance of proper dosing, duration, and medical supervision to prevent misuse and potential harm.[21]

The majority of respondents (62.2%) expressed concern about the safety of tetracycline in children, citing potential adverse events. The most commonly mentioned side effects—tooth discoloration (34.1%), nausea (25.5%), and vomiting (16.5%)—align with established risks associated with tetracycline use in children.[33,34] The concern about tooth discoloration, a well-documented risk in children under eight years of age, may be driving respondents' perceptions.[2,35,36]

While it is encouraging that respondents are aware of potential adverse effects, this highlights the need for healthcare providers to engage in open discussions with parents and caregivers about the benefits and risks of antibiotic use, including tetracycline, to ensure responsible use and minimize harm. A notable proportion of respondents (37.8%) were aware of oral adverse effects associated with tetracycline use in children, exceeding the previously reported 20%.[17] This finding highlights a significant concern among parents and caregivers about the safety of this antibiotic.

The high prevalence of tooth discoloration (90.1%) as the most common oral adverse effect aligns with previous studies, which have consistently shown that tetracycline use in children can lead to permanent tooth discoloration.[2,37,38] This is particularly alarming, as tooth discoloration can have long-lasting aesthetic and psychological impacts on children, potentially persisting into adulthood.[39,40]

Respondents exhibited varying levels of certainty regarding the safety of tetracycline, with 41.9% unsure of its safety, compared to 24.3% who were sure, and 33.7% who believed it was unsafe for pediatric use. The reason for this may stem from a lack of clear guidelines or education on tetracycline use in children. In contrast, a clear majority (63.7%) believed tetracycline is contraindicated during pregnancy, aligning with previous reports and guidelines from the American College of Obstetricians and Gynecologists (ACOG) and the Centers for Disease Control and Prevention (CDC).[33,34,41] The cautious approach to prescribing tetracycline during pregnancy is likely due to the risks of fetal harm, tooth discoloration, and the emphasis on avoiding self-medication during pregnancy.[17] Notably, an overwhelming majority (97.4%) of respondents reported never advising pregnant women to use tetracycline, indicating that healthcare providers are aware of the potential risks and are exercising caution when prescribing this antibiotic during pregnancy.

Overall, the findings emphasize the need for education and awareness campaigns to promote responsible use of antibiotics, improve knowledge about potential oral adverse effects, and reduce the risk of antibiotic resistance.

Conclusion

This study highlights poor knowledge, practice, and misuse of antibiotics and tetracycline in children, with inadequate parental/caregiver awareness of oral side effects. There is a need for education and awareness campaigns to promote responsible antibiotic use among parents and caregivers, particularly regarding tetracycline use in children. The significant knowledge gaps and misconceptions identified in this study highlight the potential for misuse and contribute to the growing concern of antibiotic resistance.

Healthcare professionals should emphasize the risks associated with tetracycline use in children, particularly the potential for permanent tooth discoloration. A multidisciplinary approach is essential to promote responsible antibiotic use, reduce resistance, and ensure the safe and effective use of antibiotics in children. Policymakers should enforce prescription requirements and regulate antibiotic sales to prevent misuse. By addressing these knowledge gaps and promoting responsible antibiotic use, we can safeguard the health and well-being of Nigerian children and contribute to a global effort to combat antibiotic resistance.

Acknowledgement: We acknowledge the parents and caregivers who participated in the study for their contribution.

Financial Disclosure: Nil.

Conflict of Interest: None declared.

Use of AI for Writing Assistance: No AI technologies utilized.

References

- Kline JM, Wietholter JP, Kline VT, Confer J. Pediatric antibiotic use: A focused review of fluoroquinolones and tetracyclines. US Pharm 2012;37(8):56–9.
- Vennila V, Madhu V, Rajesh R, Ealla KKR, Velidandla SR, Santoshi S. Tetracycline-induced discoloration of deciduous teeth: Case series. J Int Oral Health 2014;6(3):115–9.
- Dominguez RV, Acedo Diaz-pache MV, Gil MA. Prevalence of tooth discoloration from tetracycline in a school population. Av Odontostomatol 1998;5(8):541–6.
- Sede MA. Tetracycline tooth discoloration in Benin City. J Med Biomed Res 2004;3(1):53–8.
- Gbadebo SO, Ajayi DM. Self-reported tooth discolorations among patients seen at dental center university college hospital Ibadan. J West Afr Coll Surg 2015;5(3):66–77.
- Enabulele JE, Omokhua HA. Pattern of intrinsic tooth discoloration in a tertiary health facility in Nigeria. Niger J Restor Dent 2018;3:15–7.
- Itimi E, Ibhawoh LO, Enabulele JE. Satisfaction with vital tooth bleaching among a group of Nigerian patients. Sci Arch Dent Sci 2021;4:57–64.
- Shehadeh M, Suaifan G, Darwish RM, Wazaify M, Zaru L, Alja'fari S. Knowledge, attitudes, and behavior regarding antibiotic use and misuse among adults in the community of Jordan: a pilot study. Saudi Pharm J. 2012;20(2):125–33.
- 9. Shankar PR, Partha P, Shenoy N. Self-medication and non-doctor prescription practices in Pokhara Valley, Western Nepal: A questionnaire-based study. BMC Fam Pract 2002;3:17.
- Okoh M, Enabulele JE. Herbal/traditional medicine use and self-medication among patients prior to seeking oral health care in a tertiary health facility in Nigeria. J Med Res 2017;3:127–31.
- American Academy of Pediatrics. Tetracyclines. In Pickering LK, Baker CJ, Kimberlin DW, Long SS, editors. Red Book: 2009 report of the Committee on Infectious Diseases. 28th ed. Elk Grove Village: American Academy of Pediatrics; 2009.
- 12. Alfa J, Adigwe OP. Rational use of medicines in Nigeria: A critical review. J Biol Agric Healthc 2014;4(16):89–98.
- Qualtrough A, Burke F. Tooth discoloration and tetracycline. J Dent 1994;22(2):113–6.
- Okeke IN, Aboderin OA, Byarugaba DK, Ojo KK, Opintan JA. Growing problem of multidrug-resistant enteric pathogens in Africa. Emerg Infect Dis 2007;13:1640–6.
- Sapkota RE, Coker ME, Goldstein RE, Atkinson NL, Sweet JS, Sopeju PO, et al. Self-medication with antibiotics for treatment of menstrual symptom in South-West Nigeria. BMC Public Health 2010;10:1471.
- Cochran WG. Sampling techniques. 3rd ed. New York: John Wiley & Sons; 1977.
- Enabulele JE, Chukwumah NM, Enabulele O. Tetracycline use in children and knowledge of its oral implications among nursing mothers. Pediatr Dent J 2020;30:224–30.
- Okolo CC, Adeyemo YI, Oguchi CO, Malami AB, Olatosi OO, Oredugba F. An analysis of the practice of accompanying pediatric patients for dental treatment in Kano, Nigeria. Niger J Med 2022;31:581–4.

- Bankole OO, Olanloye OM, Ayebameru OE, Popoola BO. Attitude of some Nigerian parents toward their presence in the operatory during dental treatment of their children. Int J Clin Pediatr Dent 2021;14(S-2):S167–72.
- Fazli M, Kavandi M. Parents' anxiety on children cooperation at dental visit. Procedia Soc Behav Sci 2015;205:117–21.
- Asuquo FE, Akpan-Idiok PA. The exceptional role of women as primary caregivers for people living with HIV/AIDS in Nigeria, West Africa. In Suggestions for Addressing Clinical and Non-Clinical Issues in Palliative Care. London: IntechOpen; 2021.
- 22. Olatosi OO, Mgbemere OJ, Oyapero A, Omotuloye AS, Okolo CC. Awareness and preferred mode of getting information on first aid management of avulsed permanent teeth: Survey of Nigerian mothers. Pesqui Bras Odontopediatria Clin Integr 2020;11:21–4.
- Ezenwa MO, Nwagbara NV. Roles of personality traits, educational level, duration of illness in illness behavior among hypertensive patients in Anambra State, Nigeria. Niger J Psychol Res 2020;16(2):14–22.
- Latunji OO, Akinyemi OO. Factors influencing health-seeking behavior among civil servants in Ibadan, Nigeria. Ann Ib Postgrad Med 2018;16(1):52–60.
- 25. U.S. Department of Health and Human Services, Office of Disease Prevention and Health Promotion. Social determinants of health. Available at: https://odphp.health.gov/healthypeople/priority-areas/social-determinants-health. Accessed May 8, 2024.
- World Health Organization. Social determinants of health. Available at: https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1. Accessed May 8, 2024.
- 27. The Community Guide. Advancing health equity. Available at: https://www.thecommunityguide.org/pages/advancing-health-equity.html. Accessed May 8, 2024.
- Ashiru-Oredope D, Hopkins S, Vasandani S, Umoh E, Oloyede O, Nilsson A, et al; #ECDCAntibioticSurvey Project Advisory Group. Healthcare workers' knowledge, attitudes, and behaviors with respect to antibiotics, antibiotic use, and antibiotic resistance across 30 EU/EEA countries in 2019. Euro Surveill 2021;26(12):1900633.
- 29. Cambaco O, Alonso-Menendez Y, Kinsman J, Sigauque B, Wertheim H, Do N, et al. Community knowledge and practices regarding antibiotic use in rural Mozambique: Where is the starting point for prevention of antibiotic resistance? BMC Public Health 2020;20:1183.

- Mboya EA, Davies ML, Horumpende PG, Ngocho JS. Inadequate knowledge on appropriate antibiotics use among clients in the Moshi municipality Northern Tanzania. PLoS One 2020;15(9):e0239388.
- Llor C, Bjerrum L. Antimicrobial resistance: Risk associated with antibiotic overuse and initiatives to reduce the problem. Ther Adv Drug Saf 2014;5:229.
- 32. Abegaz NT, Berhe H, Gebretekle GB. Mothers/caregivers' healthcare-seeking behavior toward childhood illness in selected health centers in Addis Ababa, Ethiopia: A facility-based cross-sectional study. BMC Pediatr 2019;19:220.
- Shutter MC, Akhondi H. Tetracycline. In StatPearls. Treasure Island: StatPearls Publishing; 2024.
- Ramesh SA, Kabde PR, Chavale GV, Sapkal HS. A review on contraindications of tetracycline in pregnancy. World J Pharm Life Sci 2024;10(7):317–23.
- Sánchez AR, Rogers RS, Sheridan PJ. Tetracycline and other tetracycline-derivative staining of the teeth and oral cavity. Int J Dermatol 2004;43(10):709–15.
- Conchie JM, Munroe JD, Anderson DO. The incidence of staining of permanent teeth by the tetracyclines. Can Med Assoc J 1970;103(4):351–6.
- 37. American Academy of Pediatrics. Tetracyclines. In Kimberlin DW, Jackson MA, Long SS, editors. Red Book: 2018 Report of the Committee on Infectious Diseases. 31st ed. Itasca: American Academy of Pediatrics; 2018.
- Zhu Z, Yu Q, Qi G, Yang J, Ni Y, Ruan W, et al. Tigecycline-induced tooth discoloration in children younger than eight years. Antimicrob Agents Chemother 2021;65(9):e0085421.
- Ibiyemi O, Taiwo JO. Psychosocial aspect of anterior tooth discoloration among adolescents in Igbo-Ora, Southwestern Nigeria. Ann Ib Postgrad Med 2011;9(2):94–9.
- 40. Ogunsuji OO, Molokwu CF, Oyedeji KE, Taiwo IM, Udoji CS, Adegoke KT, et al. Psychosocial impact of tooth discoloration among 12- to 15-year-old school children in Ibadan Northwest Local Government Area of Oyo State, Nigeria. Oral Health Dent Sci 2021;5(3):1–7.
- 41. Yu PA, Tran EL, Parker CM, Kim HJ, Yee EL, Smith PW, et al. Safety of antimicrobials during pregnancy: A systematic review of antimicrobials considered for treatment and postexposure prophylaxis of plague. Clin Infect Dis 2020;70(Suppl 1):S37–50.